

COMMON QUARTERLY EXAMINATION-2024-25

Time Allowed : 3.00 Hours]

MATHEMATICS

[Max. Marks : 100

PART - I

14x1=14

1. Choose the best answer:
 1. If there are 1024 relations from a set $A = \{1, 2, 3, 4, 5\}$ to a set B . Then the number of elements in B is
 - a) 3 b) 2 c) 4 d) 8
 2. $f(x) = (x+1)^3 - (x-1)^3$ represents a function which is
 - a) linear b) cubic c) reciprocal d) quadratic
 3. Euclid's division lemma states that for positive integers a and b , there exists unique integers q and r such that $a=bq+r$, where r must satisfy.
 - a) $1 < r < b$ b) $0 < r < b$ c) $0 \leq r < b$ d) $0 < r \leq b$
 4. If $A = 2^{65}$ and $B = 2^{64} + 2^{63} + 2^{62} + \dots + 2^0$. which of the following is true?
 - a) B is 2^{64} more than A b) A and B are equal
 - c) B is larger than A by 1 d) A is larger than B by 1
 5. A system of three linear equations in three variables is inconsistent if their planes.
 - a) Intersect only at a points b) Intersect in a line
 - c) Coincides with each other d) Do not intersect
 6. $y^2 + \frac{1}{y^2}$ is not equal to
 - a) $\frac{y^4 + 1}{y^2}$ b) $\left(y + \frac{1}{y}\right)^2$ c) $\left(y - \frac{1}{y}\right)^2 + 2$ d) $\left(y + \frac{1}{y}\right)^2 - 2$
 7. Graph of a quadratic equation is a -----
 - a) Straight line b) Circle c) Parabola d) Hyperbola
 8. If in triangle ABC and DEF , $\frac{AB}{DE} = \frac{BC}{FD}$, then they will be similar, when
 - a) $\angle B = \angle E$ b) $\angle A = \angle D$ c) $\angle B = \angle D$ d) $\angle A = \angle F$
 9. If in $\triangle ABC$, $DE \parallel BC$, $AB = 3.6$ cm, $AC = 2.4$ cm and $AD = 2.1$ cm then the length of AE is
 - a) 1.4 cm b) 1.8 cm c) 1.2 cm d) 1.05 cm
 10. The straight line given by the equation $x = 11$ is
 - a) Parallel to X axis b) Parallel to Y axis
 - c) Passing through the origin d) Passing through the point $(0, 11)$
 11. If slope of the line PQ is $\frac{1}{\sqrt{3}}$ then slope of the perpendicular bisector of PQ is
 - a) $\sqrt{3}$ b) $-\sqrt{3}$ c) $\frac{1}{\sqrt{3}}$ d) 0
 12. Consider four straight lines: (i) $l_1: 3y=4x+5$ (ii) $l_2: 4y=3x-1$ (iii) $l_3: 4y+3x=7$ (iv) $l_4: 4x+3y=2$
Which of the following statement is true?
 - a) l_1 and l_2 are perpendicular b) l_1 and l_4 are parallel
 - c) l_2 and l_4 are perpendicular d) l_2 and l_3 are parallel
 13. The value of $\sin^2 \theta + \frac{1}{1 + \tan^2 \theta}$ is equal to
 - a) $\tan^2 \theta$ b) 1 c) $\cot^2 \theta$ d) 0
 14. If $5x = \sec \theta$ and $\frac{5}{x} = \tan \theta$, then $x^2 - \frac{1}{x^2}$ is equal to
 - a) 25 b) $\frac{1}{25}$ c) 5 d) 1

PART - II

Answer any 10 questions. Question No. 28 is compulsory.

10x2=20

15. If $B \times A = \{(-2, 3), (-2, 4), (0, 3), (0, 4), (3, 3), (3, 4)\}$ find A and B .
16. Represent the function $f = \{(1, 2), (2, 2), (3, 2), (4, 3), (5, 4)\}$ through (i) an arrow diagram (ii) a table form
17. Is $7 \times 5 \times 3 \times 2 + 3$ a composite number? Justify your answer.
18. Find the number of terms in the G.P. 4, 8, 16, 8192?

19. Find the LCM of $8x^4y^2$, $48x^2y^4$.
20. Solve the equation $4x^2 - 7x - 2 = 0$ by factorization method.
21. If $\triangle ABC$ is similar to $\triangle DEF$ such that $BC = 3$ cm, $EF = 4$ cm and area of $\triangle ABC = 54\text{cm}^2$. Find the area of $\triangle DEF$.
22. State the angle bisector theorem.
23. Show that the points $P(-1.5, 3)$, $Q(6, -2)$, $R(-3, 4)$ are collinear.
24. Show that the straight lines $x - 2y + 3 = 0$ and $6x + 3y + 8 = 0$ are perpendicular.
25. Find the equation of a line whose intercepts on the x and y axes are 4 and -6.
26. Prove that $\sqrt{\frac{1 + \cos\theta}{1 - \cos\theta}} = \operatorname{cosec}\theta + \cot\theta$
27. Find the sum : $1 + 2 + 3 + \dots + 60$.
28. If the difference between a number and its reciprocal is $24/5$, find the number.

PART - III

Answer the following any 10 questions. Q.No.42 is compulsory.

10x5=50

29. Let $A = \{x \in W \mid x < 2\}$, $B = \{x \in N \mid 1 < x \leq 4\}$ and $C = \{3, 5\}$. Verify that $A \times (B \cup C) = (A \times B) \cup (A \times C)$.
30. If the function $f: R \rightarrow R$ is defined by $f(x) = \begin{cases} 2x + 7; & x < -2 \\ x^2 - 2; & -2 \leq x < 3 \\ 3x - 2; & x \geq 3 \end{cases}$ find the values of
 - i) $f(4)$
 - ii) $f(-2)$
 - iii) $f(4) + 2f(1)$
 - iv) $\frac{f(1) - 3f(4)}{f(-3)}$
31. Rekha has 15 square colour papers of sizes 10 cm, 11 cm, 12 cm.....24cm. How much area can be decorated with these colour papers?
32. Find the GCD of $6x^3 - 30x^2 + 60x - 48$ and $3x^3 - 12x^2 + 21x - 18$.
33. If $9x^4 + 12x^3 + 28x^2 + ax + b$ is a perfect square, find the values of a and b.
34. If $A = \frac{2x+1}{2x-1}$, $B = \frac{2x-1}{2x+1}$ find $\frac{1}{A-B} - \frac{2B}{A^2-B^2}$
35. State and prove Basic Proportionality theorem.
36. A boy height 90 cm is walking away from the base of a lamp post at a speed of 1.2m/sec. If the lamp post is 3.6 m above the ground. Find the length of his shadow cast after 4 seconds.
37. Find the Area of the quadrilateral whose vertices are at $(-9, 0)$, $(-8, 6)$, $(-1, -2)$ and $(-6, -3)$
38. Let $A(3, -4)$, $B(9, -4)$, $C(5, -7)$ and $D(7, -7)$. Show that ABCD is a trapezium.
39. A line makes positive intercepts on coordinate axes whose sum is 7 and it passes through $(-3, 8)$. Find its equation.
40. Prove that $\sin^2 A \cos^2 B + \cos^2 A \sin^2 B + \cos^2 A \cos^2 B + \sin^2 A \sin^2 B = 1$.
41. If $\frac{\cos\alpha}{\sin\beta} = m$ and $\frac{\cos\alpha}{\sin\beta} = n$, then prove that $(m^2 + n^2) \cos^2 \beta = n^2$
42. If $S_1, S_2, S_3, \dots, S_m$ are the sum of n terms of m A.P.'s. Whose first terms are 1, 2, 3,m and whose common difference are 1, 3, 5, $(2m-1)$ respectively. Then show that $S_1 + S_2 + S_3 + \dots + S_m = \frac{1}{2} mn (mn+1)$.

PART - IV

Answer all of the following.

2x8=16

43. a) Construct a triangle similar to a given triangle PQR with its sides equal to $7/3$ of the corresponding sides of the triangle PQR (scale factor $7/3 > 1$) (OR)
- b) Construct a $\triangle PQR$ in which $PQ = 8$ cm, $\angle R = 60^\circ$ and the median RG from R to PQ is 5.8 cm. Find the length of the altitude from R to PQ.
44. a) Graph the following linear function $y = \frac{1}{2}x$. Identify the constant of variation and verify it with the graph. Also (i) Find y when $x = 9$, (ii) Find x when $y = 7.5$. (OR)
- b) Nishanth is the winner in a Marathon race of 12 km distance. He ran at the uniform speed of 12 km/hr and reached the destination in 1 hour. He was followed by Aradhana, Jayanth, Sathya and Swetha with their respective speed of 6 km/hr, 4 km/hr, 3 km/hr and 2 km/hr and they covered the distance in 2 hrs, 3 hrs, 4 hrs and 6 hours respectively. Draw the speed - time graph and use it to find the time taken to Kanchik with his speed of 2.4 km/hr.

KK / 10 / Mat / 2

MATHEMATICS

Part – I

Q.No.	Option	Answer
1.	b	2
2.	d	quadratic
3.	c	$0 \leq n \leq 6$
4.	d	A is larger than B by 1.
5.	d	Do not intersect.
6.	b.	$(y + \frac{1}{y})^2$
7.	c.	Parabola.
8.	c	$\angle B = \angle D$
9.	a	1.4 cm.
10.	b.	Parallel to Y-axis.
11.	b	$-\sqrt{3}$
12.	c	l_2 and l_4 are perpendicular.
13.	b	1.
14.	b	$\frac{1}{25}$.

Part - II

15. Solution:-

Given: $B \times A = \{(-2, 3), (-2, 4), (0, 3), (0, 4), (3, 3), (3, 4)\}$.

B = Set of all first co ordinates of element of $B \times A$.

$$B = \{-2, 0, 3\}$$


A = set of all second co ordinates of element of $B \times A$.

$$A = \{3, 4\}$$

Hence, $A = \{3, 4\}$, $B = \{-2, 0, 3\}$.

16. Solution:-

i] AN ARROW DIAGRAM.



ii] A TABLE FORM.

x	1	2	3	4	5
$f(x)$	2	2	2	3	4

17. Solution:-

Yes, the given number is a composite number, because.

$$7 \times 5 \times 3 \times 2 + 3 = 3 \times (7 \times 5 \times 2 + 1) = 3 \times 71$$

Since, the given number can be factorized in terms of two primes, it is a composite number.

18. Solution:-

The Given G.P is 4, 8, 16, ..., 8192.

$$r = \frac{t_2}{t_1} = \frac{8}{4} = 2 \Rightarrow \boxed{r=2}, \boxed{a=4}$$

Let the n^{th} term be 8192.

$$t_n = 8192$$

$$t_n = ar^{n-1}$$

$$8192 = 4(2)^{n-1}$$

$$\frac{8192}{4} = 2^{n-1}$$

$$2^{n-1} = 2048$$

$$2^{n-1} = 2^{11}$$

Since, the base are equal, Powers are also equal,

$$n-1 = 11$$

Hence, $n = 12$

There are 12 terms in the given sequence.

19. Solution:-

To FIND: LCM of $8x^4y^2$, $48x^2y^4$.

$$8x^4y^2 = 2 \times 2 \times 2 \times x^2 \times x^2 \times y^2$$

$$48x^2y^4 = 2 \times 2 \times 2 \times 2 \times 3 \times x^2 \times y^2 \times y^2$$

$\therefore$ LCM of $8x^4y^2$, $48x^2y^4$ is $8 \times 6 \times x^2 \times x^2 \times y^2 \times y^2$.

$$\Rightarrow 48x^4y^4$$

Hence,

$\therefore$ LCM of $8x^4y^2$, $48x^2y^4$ is $48x^4y^4$.

20. Solution:-

by using factorization method,

$$4x^2 - 7x - 2 = 0. \quad -x' - 8$$

$$4x^2 - 8x + x - 2 = 0$$

$$4x[x-2] + [x-2] = 0. \quad 1 \mid -8$$

$$(x-2)(4x+1) = 0. \quad -7$$

$$x-2=0 \quad 4x+1=0$$

$$x=2 \quad x=-\frac{1}{4}$$

Hence,

The roots are 2 and $-\frac{1}{4}$.

21. Solution:-

Since, the ratio of area of two similar triangles is equal to the ratio of the squares of any two corresponding sides,

$$\text{we have, } \frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DEF)} = \frac{BC^2}{EF^2}$$

$$\Rightarrow \frac{54}{\text{Area } \triangle DEF} = \frac{3^2}{4^2}$$

Hence,


$$\text{Area } \triangle DEF = \frac{54 \times 16}{9} = 96 \text{ cm}^2$$

22. Solution:-

STATEMENT: Angle Bisector Theorem.

The internal bisector of an angle of a triangle divides the opposite side internally in the ratio of the corresponding sides containing the angle.

DIAGRAM:



23. Solution:-

The points are $P(-1.5, 3)$, $Q(6, -2)$
 $R(-3, 4)$
 x_1, y_1 x_2, y_2 x_3, y_3

$$\begin{aligned} \text{Area of } \triangle PQR &= \frac{1}{2} \{ (x_1 y_2 + x_2 y_3 + x_3 y_1) \\ &\quad - (x_2 y_1 + x_3 y_2 + x_1 y_3) \} \\ &\quad \text{Sq. units} \\ &= \frac{1}{2} \{ (3 + 24 - 9) - (18 + 6 - 6) \} \\ &= \frac{1}{2} \{ (18 - 18) \} = 0 \end{aligned}$$

Hence,

$\therefore$ they given points are collinear.

24. Solution:-

slope of st. line $x - 2y + 3 = 0$ is

$$m_1 = \frac{-\text{coeff. of } x}{\text{coeff. of } y} = \frac{-1}{-2} = \frac{1}{2}$$

slope of st. line $6x + 3y + 8 = 0$ is.

$$m_2 = \frac{-6}{3} = -2$$

Now,

$$m_1 \times m_2 = -1$$

$$m_1 \times m_2 = \frac{1}{2} \times -2 = -1$$

Hence,

the two st. lines are perpendicular.

25. Solution:-

Given, x -intercept $\Rightarrow a = 4$

y -intercept $\Rightarrow b = -6$.

Intercepts form,

$$\frac{x}{a} + \frac{y}{b} = 1$$

$$\frac{x}{4} + \frac{y}{-6} = 1$$

$$\frac{-6x + 4y}{-24} = 1$$

$$-6x + 4y = -24$$

$\times$ by -1

$$6x - 4y = 24$$

$$\Rightarrow 3x - 2y = 12$$

Hence,

they required equation is

$$3x - 2y = 12$$

26. Solution:-

$$\sqrt{\frac{1+\cos\theta}{1-\cos\theta}} = \sqrt{\frac{1+\cos\theta}{1-\cos\theta} \times \frac{1+\cos\theta}{1+\cos\theta}}$$

$$= \sqrt{\frac{(1+\cos\theta)^2}{1-\cos^2\theta}} = \sqrt{\frac{(1+\cos\theta)^2}{\sin^2\theta}}$$

$$= \sqrt{\left(\frac{1+\cos\theta}{\sin\theta}\right)^2} = \frac{1+\cos\theta}{\sin\theta}$$

$$= \frac{1}{\sin\theta} + \frac{\cos\theta}{\sin\theta}$$

$$= \csc\theta + \cot\theta$$

Hence proved.

27. Solution:-

$$1+2+3+\dots+n = \frac{n(n+1)}{2}$$

$$n=60$$

$$1+2+3+\dots+60 = \frac{60 \times 61}{2} = 30 \times 61 = 1830$$

Hence,

$$1+2+3+\dots+60 = 1830 \text{ ✓}$$

28. Solution:-

Let the number be x &
its reciprocal is $\frac{1}{x}$.

$$\therefore x - \frac{1}{x} = \frac{24}{5}$$

$$\frac{x^2-1}{x} = \frac{24}{5}$$

$$5(x^2-1) = 24x$$

$$5x^2 - 24x - 5 = 0$$

$$(x-5)(5x+1) = 0$$

$$x=5, x=-\frac{1}{5}$$

Hence, The required numbers are 5 & $-\frac{1}{5}$.

Part - III

29. Solution:-

Given that,

$$A = \{x \in W \mid x < 2\} = \{0, 1\}$$

$$B = \{x \in N \mid 1 < x \leq 4\} = \{2, 3, 4\}$$

$$C = \{3, 5\}$$

$$B \cup C = \{2, 3, 4\} \cup \{3, 5\} = \{2, 3, 4, 5\}$$

$$A \times (B \cup C) = \{0, 1\} \times \{2, 3, 4, 5\}$$

$$A \times (B \cup C) = \{(0, 2), (0, 3), (0, 4), (0, 5), (1, 2), (1, 3), (1, 4), (1, 5)\}$$

→ ①

$$A \times B = \{0, 1\} \times \{2, 3, 4\}$$

$$= \{(0, 2), (0, 3), (0, 4), (1, 2), (1, 3), (1, 4)\}$$

$$A \times C = \{0, 1\} \times \{3, 5\}$$

$$= \{(0, 3), (0, 5), (1, 3), (1, 5)\}$$

$$(A \times B) \cup (A \times C)$$

$$= \{(0, 2), (0, 3), (0, 4), (1, 2), (1, 3), (1, 4)\} \cup \{(0, 3), (0, 5), (1, 3), (1, 5)\}$$

$$= \{(0, 2), (0, 3), (0, 4), (0, 5), (1, 2), (1, 3), (1, 4), (1, 5)\}$$

From ① & ②, we get.

→ ②

$$A \times (B \cup C) = (A \times B) \cup (A \times C)$$

Hence Proved ✓

30. Solution:-

Given function $f: R \rightarrow R$ is defined

$$\text{by } f(x) = \begin{cases} 2x+7 & ; x < -2 \\ x^2-2 & ; -2 \leq x < 3 \\ 3x-2 & ; x \geq 3 \end{cases}$$

i] $f(4) \Rightarrow$ lie third interval $x=4$.

$$f(x) = 3x-2$$

$$f(4) = 3(4)-2 = 12-2 = 10$$

ii]. $f(-2) \Rightarrow$ lie second interval $x=-2$.

$$f(x) = x^2-2$$

$$f(-2) = (-2)^2-2 = 4-2 = 2$$

$$\text{iii]. } f(4) + 2f(1) = 10 + 2(1^2-2)$$

$$= 10 + 2(1-2)$$

$$= 10 + 2(-1) = 10-2$$

$$= 8$$

$$\text{iv]. } \frac{f(1) - 3f(4)}{f(-3)} = \frac{-1 - 3(10)}{2(-3)+7} = \frac{-1-30}{-6+7} = \frac{-31}{1}$$

$$= -31 \text{ ✓}$$

Hence.

$$\text{i] } f(4)=10, \text{ ii] } f(-2)=2 \text{ iii] } f(4)+2f(1)=8$$

$$\text{iv]. } \frac{f(1) - 3f(4)}{f(-3)} = -31 \text{ ✓}$$

31. Solution:-

Total area can be decorated } = $10^2 + 11^2 + 12^2 + \dots + 24^2$

$$= (1^2 + 2^2 + \dots + 24^2) - (1^2 + 2^2 + \dots + 9^2)$$

$$\left[\therefore 1^2 + 2^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6} \right]$$

$$= \frac{24 \times 25 \times 49}{6} - \frac{9 \times 10 \times 19}{6}$$

$$= \frac{12^4}{2 \times 3} - \frac{3 \times 10^5}{1 \times 2}$$

$$= 4900 - 285$$

$$= 4615 \text{ cm}^2$$

Hence

Rekha has covered, area can be decorated with these colour papers = 4615 cm^2 h.

32. Solution:-

Let $f(x) = 6x^3 - 30x^2 + 60x - 48$
 $= 6(x^3 - 5x^2 + 10x - 8)$ and.

$g(x) = 3x^3 - 12x^2 + 21x - 18$
 $= 3(x^3 - 4x^2 + 7x - 6)$

$$\begin{array}{r} x^3 - 5x^2 + 10x - 8 \\ x^3 - 4x^2 + 7x - 6 \quad (-) \\ \hline x^2 - 3x + 2 \end{array}$$

$$\begin{array}{r} x^2 - 3x + 2 \\ x^3 - 5x^2 + 10x - 8 \quad (-) \\ \hline x^3 - 3x^2 + 2x \quad (-) \\ \hline -2x^2 + 8x - 8 \quad (-) \\ \hline -2x^2 + 6x - 4 \quad (-) \\ \hline 2x - 4 \quad (-) \\ \hline 2x - 4 \quad (-) \\ \hline 0 \end{array} = 2(x-2)$$

$$\begin{array}{r} x-1 \\ x^2 - 3x + 2 \\ \hline x^2 - 2x \quad (-) \\ \hline -x + 2 \quad (-) \\ \hline (-) - x + 2 \quad (-) \\ \hline 0 \quad \text{remainder as zero.} \end{array}$$

GCD of leading co-eff's 3 & 6 is 3.

Hence,

$$\text{GCD} [(6x^3 - 30x^2 + 60x - 48, 3x^3 - 12x^2 + 21x - 18)] = 3(x-2)$$

33. Solution:-

$$\begin{array}{r} 3x^2 + 2x + 4 \\ 3x^2 \quad 9x^4 + 12x^3 + 28x^2 + 9x + 6 \quad (-) \\ \hline 6x^2 + 2x \quad 12x^3 + 28x^2 \quad (-) \\ \hline 6x^2 + 4x + 4 \quad 12x^3 + 4x^2 \quad (-) \\ \hline 6x^2 + 4x + 4 \quad 24x^2 + 9x + 6 \quad (-) \\ \hline 24x^2 + 16x + 16 \quad (-) \\ \hline 0 \end{array}$$

Because, the given polynomial is a perfect square.

Equating x -co-efficients, we get.

$$a = 16$$

Equating constant term, we get.

$$b = 16$$

Hence,

$$a = 16, b = 16$$

34. Solution:-

Given, $A = \frac{2x+1}{2x-1}$, $B = \frac{2x-1}{2x+1}$.

$$\begin{aligned} \frac{1}{A-B} - \frac{2B}{A^2-B^2} &= \frac{1}{A-B} - \frac{2B}{(A+B)(A-B)} \\ &= \frac{1}{A-B} \left[1 - \frac{2B}{A+B} \right] \\ &= \frac{1}{A-B} \left[\frac{A+B-2B}{A+B} \right] \\ &= \frac{1}{A-B} \left[\frac{A-B}{A+B} \right] \\ &= \frac{1}{A+B} = \frac{1}{\left(\frac{2x+1}{2x-1}\right) + \left(\frac{2x-1}{2x+1}\right)} \\ &= \frac{1}{\frac{(2x+1)^2 + (2x-1)^2}{(2x)^2 - 1^2}} \\ &= \frac{4x^2 - 1}{4x^2 + 1 + 4x + 4x^2 + 1 - 4x} \\ &= \frac{4x^2 - 1}{8x^2 + 2} \\ &= \frac{4x^2 - 1}{2(4x^2 + 1)} \end{aligned}$$

Hence,

$$\boxed{\frac{1}{A-B} - \frac{2B}{A^2-B^2} = \frac{4x^2-1}{2(4x^2+1)} \text{ Proved.}}$$

35. Solution:-

Basic Proportionality Theorem (or) Thales Theorem


STATEMENT:-

A st. line drawn parallel to a side of triangle intersecting the other two sides, divides the sides in the same ratio.

GIVEN:-

In $\triangle ABC$,

D is a point on AB
E is a point on AC.



TO PROVE:-

$$\frac{AD}{DB} = \frac{AE}{EC}$$

CONSTRUCTION:

Draw a line $DE \parallel BC$.

$\angle ABC = \angle ADE = \angle 1$ (corresponding angles are equal, $DE \parallel BC$)

$\angle ACB = \angle AED = \angle 2$ (corresponding angles are equal, $DE \parallel BC$)

$\angle DAE = \angle BAC = \angle 3$ (Both triangles have a common angle).

$\triangle ABC \sim \triangle ADE$ by AAA similarity.

$\frac{AB}{AD} = \frac{AC}{AE}$ [corresponding sides are proportional.]

$\frac{AD+DB}{AD} = \frac{AE+EC}{AE}$ [Split AB & AC using the point D & E]


$1 + \frac{DB}{AD} = 1 + \frac{EC}{AE}$ on simplification.

$\frac{DB}{AD} = \frac{EC}{AE}$ Cancelling 1 on both sides.

$\frac{AD}{DB} = \frac{AE}{EC}$ Taking reciprocals.

Hence Proved.

36. Solution:-



Given, Speed = 1.2 m/s ,
 time = 4 seconds.
 distance = speed $\times$ time.
 $= 1.2 \times 4 = 4.8 \text{ m}$.

distance = 4.8 m .

Let x be the length of the shadow after 4 seconds.

Since, $\triangle ABE \sim \triangle CDE$,

$$\frac{BE}{DE} = \frac{AB}{CD} \quad [90 \text{ cm} = 0.9 \text{ m}].$$

$$\frac{4.8+x}{x} = \frac{3.6}{0.9}$$

$$\frac{4.8+x}{x} = 4.$$

$$4.8+x = 4x.$$

$$\Rightarrow 3x = 4.8$$

$$x = 1.6 \text{ m}.$$

Hence,


The length of his shadow DE = 1.6 m .

37. Solution:-

* Plot the points on the plane & take them in a counter clockwise direction.

* Let the vertices are

$A(-8, 6)$ $B(-9, 0)$
 $C(-6, -3)$ $D(-1, -2)$.



$$\begin{aligned} \text{Area of a quadrilateral ABCD} &= \frac{1}{2} \{ (x_1 - x_3)(y_2 - y_4) - (x_2 - x_4)(y_1 - y_3) \} \\ &= \frac{1}{2} \{ (-8+6)(0+2) - (-9+1)(6+3) \} \\ &= \frac{1}{2} [(-2)(2) - (-8)(9)] \\ &= \frac{1}{2} [-4 + 72] = \frac{1}{2} [68] = 34. \end{aligned}$$

Hence,

Area of a quadrilateral ABCD = 34 sq. units .

38. Solution:-

by using $m = \frac{y_2 - y_1}{x_2 - x_1}$.

Given: $A(3, -4)$ $B(9, -4)$ $C(5, -7)$ $D(7, -7)$

$$\text{Slope of AB} = \frac{-4+4}{9-3} = \frac{0}{6} = 0.$$

$$\boxed{\text{Slope of AB} = 0} \rightarrow \textcircled{1}$$

$$\text{Slope of BC} = \frac{-7+4}{5-9} = \frac{-3}{-4} = \frac{3}{4}.$$

$$\boxed{\text{Slope of BC} = \frac{3}{4}} \rightarrow \textcircled{2}$$

$$\text{Slope of CD} = \frac{-7+7}{7-5} = \frac{0}{2} = 0.$$

$$\boxed{\text{Slope of CD} = 0} \rightarrow \textcircled{3}$$

$$\text{Slope of AD} = \frac{-7+4}{7-3} = \frac{-3}{4} = -\frac{3}{4}.$$

$$\boxed{\text{Slope of AD} = -\frac{3}{4}} \rightarrow \textcircled{4}$$

From $\textcircled{1}$, $\textcircled{2}$, $\textcircled{3}$ & $\textcircled{4}$, we get,

slope of AB = slope of CD.

Since, one pair of opposite sides are parallel,

Hence, $\therefore$ ABCD is trapezium.

39. Solution:-

If a & b are the intercepts then $a+b=7$

$$\Rightarrow \boxed{b=7-a}$$

By intercept form, $\frac{x}{a} + \frac{y}{b} = 1$,

$$\text{we have, } \frac{x}{a} + \frac{y}{7-a} = 1. \rightarrow \textcircled{1}$$

As this line passes through the point $(-3, 8)$,

$$\textcircled{1} \Rightarrow \frac{-3}{a} + \frac{8}{7-a} = 1$$

$$\Rightarrow -3(7-a) + 8a = a(7-a)$$

$$-21 + 3a + 8a = 7a - a^2$$

$$\Rightarrow a^2 + 4a - 21 = 0$$

$$a^2 - 3a + 7a - 21 = 0$$

$$a(a-3) + 7(a-3) = 0$$

$$(a-3)(a+7) = 0$$

$$a = 3 \text{ (or)} a = -7$$

Put $a = 3$, $b = 7 - a = 7 - 3 = 4 \Rightarrow b = 4$

Put $a = -7$, $b = 7 - (-7) = 7 + 7 = 14 \Rightarrow b = 14$

Since, a is positive,

So, $a = 3$ & $b = 4$

Hence $\frac{x}{3} + \frac{y}{4} = 1$

$\therefore 4x + 3y = 12 \Rightarrow 4x + 3y - 12 = 0$
is the required equation.

40. Solution:-

$$\text{L.H.S} = \sin^2 A \cos^2 B + \cos^2 A \sin^2 B + \cos^2 A \cos^2 B + \sin^2 A \sin^2 B$$

$$= \sin^2 A \cos^2 B + \sin^2 A \sin^2 B + \cos^2 A \sin^2 B + \cos^2 A \cos^2 B$$

$$= \sin^2 A [\cos^2 B + \sin^2 B] + \cos^2 A [\sin^2 B + \cos^2 B]$$

[since, $\sin^2 B + \cos^2 B = 1$]

$$= \sin^2 A [1] + \cos^2 A [1]$$

$$= \sin^2 A + \cos^2 A$$

$$= 1 \quad [\because \sin^2 A + \cos^2 A = 1]$$

$$= \text{R.H.S.}$$

$$\therefore \text{L.H.S} = \text{R.H.S.}$$

$$\therefore \sin^2 A \cos^2 B + \cos^2 A \sin^2 B + \cos^2 A \cos^2 B + \sin^2 A \sin^2 B = 1$$

41. Solution:-

Given that,

$$\frac{\cos \alpha}{\cos \beta} = m \quad \bigg/ \quad \frac{\cos \alpha}{\sin \beta} = n$$

[Question Paper mistake in this sum]

$$\therefore m^2 + n^2 = \left[\frac{\cos \alpha}{\cos \beta} \right]^2 + \left[\frac{\cos \alpha}{\sin \beta} \right]^2$$

$$= \frac{\cos^2 \alpha}{\cos^2 \beta} + \frac{\cos^2 \alpha}{\sin^2 \beta}$$

$$= \cos^2 \alpha \left[\frac{1}{\cos^2 \beta} + \frac{1}{\sin^2 \beta} \right]$$

$$= \cos^2 \alpha \left[\frac{\cos^2 \beta + \sin^2 \beta}{\cos^2 \beta \sin^2 \beta} \right]$$

$$= \cos^2 \alpha \left[\frac{1}{\cos^2 \beta \sin^2 \beta} \right]$$

$$= \frac{1}{\cos^2 \beta} \left[\frac{\cos^2 \alpha}{\sin^2 \beta} \right]$$

$$m^2 + n^2 = \frac{1}{\cos^2 \beta} [n^2]$$

$$\boxed{(m^2 + n^2) \cos^2 \beta = n^2}$$

Hence Proved.

42. Solution:-

S_1 denotes the sum of n terms,

$$a=1, d=1.$$

$$S_1 = \frac{n}{2} [2(1) + (n-1)(1)].$$

S_2 denotes the sum of n terms,

$$a=2, d=3.$$

$$S_2 = \frac{n}{2} [2(2) + (n-1)3].$$

S_3 denotes the sum of n terms,

$$a=3, d=5.$$

$$S_3 = \frac{n}{2} [2(3) + (n-1)5].$$

...

S_m denotes the sum of n terms,

$$a=m, d=2m-1.$$

$$S_m = \frac{n}{2} [2(m) + (n-1)(2m-1)].$$

$$\therefore S_1 + S_2 + \dots + S_m$$

$$= \frac{n}{2} [2(1+2+\dots+m) + (n-1)(1+3+5+\dots+(2m-1))]$$

$$= \frac{n}{2} \left[2 \frac{m(m+1)}{2} + (n-1) \frac{m}{2} [1+(2m-1)] \right]$$

$$a=1, d=2m-1$$

$$n=m.$$

$$= \frac{n}{2} n \left[(m+1) + \frac{(n-1)(2m)}{2} \right]$$

$$= \frac{mn}{2} \left[m+1 + \frac{2mn-2m}{2} \right]$$

$$= \frac{mn}{2} \left[\frac{2m+2+2mn-2m}{2} \right]$$

$$= \frac{mn}{2} \left[\frac{2[mn+1]}{2} \right]$$

$$= \frac{mn}{2} [mn+1].$$

Hence,

$$\therefore S_1 + S_2 + S_3 + \dots + S_m = \frac{mn}{2} [mn+1]$$



h.

Part - IV

43.a) Solution:-

Given triangle PQR.

[scale factor $\frac{7}{3} > 1$].




Construction:-

- 1]. Construct a $\triangle PQR$ with any measurement.
- 2]. Draw a ray QX making an acute angle with QR on the side opposite to vertex P .
- 3]. Locate 7 points [the greater of $7 \times \frac{7}{3}$]
 $Q_1, Q_2, Q_3, Q_4, Q_5, Q_6$ and Q_7 on QX ,
 $Q_1Q_2 = Q_2Q_3 = Q_3Q_4 = Q_4Q_5 = Q_5Q_6 = Q_6Q_7$.
- 4]. Join Q_3 to R and draw a line through Q_7 $\parallel$ to Q_3R , intersecting the extended line segment QR at R' .
- 5]. Draw line through R' Parallel to

the line RP intersecting the extending line segment QP at P' .

Then, $\triangle P'QR'$ is the required triangle each of whose sides is seven-third of the corresponding sides of $\triangle PQR$.

43) b). Solution:-




- Step 1: Draw a line segment $PQ = 8\text{cm}$
- Step 2: At P , draw PF such that $\angle QPF = 60^\circ$
- Step 3: At P , draw PF such that $\angle FPF = 90^\circ$
- Step 4: Draw the perpendicular bisector to PQ , which intersects PF at O and PQ at G .
- Step 5: With O as centre and OP as radius draw a circle.
- Step 6: From G mark arcs of radius 5.8cm on the circle. Mark them as R and S .
- Step 7: Join PH and RQ . Then $\triangle PQR$ is the required triangle.
- Step 8: From R draw a line RN perpendicular to EQ . EQ meets RN at M .
- Step 9: The length of the altitude is $RM = 3.8\text{cm}$.

44)
a)

Let us form a table for the given function

x	2	4	6	8	10	12	14	16
y	1	2	3	4	5	6	7	8

From the table we observe that as x increases, y also increases. Therefore, the variation is a direct variation. Hence, the constant of variation $k = \frac{1}{2}$



Plot the points (2,1), (4,2), (6,3), (8,4), (10,5), (12,6), (14,7) and (16,8) on the graph

We observe that the relation $y = \frac{1}{2}x$ forms a straight line.

From the graph,

- (i) when $x = 9$, we have $y = 4.5$
- (ii) when $y = 7.5$, we have $x = 15$

44)
b)

Let us form the table with the given details.

Speed x (km/hr)	12	6	4	3	2
Time y (hours)	1	2	3	4	6

From the table, we observe that as x decreases, y increases. Hence, the type is inverse variation.

$$\text{Let } y = \frac{k}{x}$$


$\Rightarrow xy = k, k > 0$ is called the constant of variation.

$$\text{From the table } k = 12 \times 1 = 6 \times 2 = \dots = 2 \times 6 = 12$$

Therefore, $xy = 12$.

Plot the points (12,1), (6,2), (4,3), (3,4), (2,6) and join these points by a smooth curve (Rectangular Hyperbola).

From the graph, we observe that Kaushik takes 5 hrs with a speed of 2.4 km/hr.



...ALL THE BEST...

Prepared By,

M.S.Mahendira Rajan, M.Sc., M.Phil., B.Ed.,

Vice Principal & Dept. In Mathematics.

Vanmathi Matric Hr.Sec. School

Vadakkanandal, Chinnasalem(Tk), Kallakurichi(Dt).

contact Number: 9500499292.